

# Intercomparison and Evaluation of Cumulus Parameterizations under Summertime Midlatitude Continental Conditions

Shaocheng Xie<sup>1\*</sup>, Kuan-Man Xu<sup>2</sup>, Richard T. Cederwall<sup>1</sup>, Peter Bechtold<sup>3</sup>,  
Anthony D. Del Genio<sup>4</sup>, Stephen A. Klein<sup>5</sup>, Douglas G. Cripe<sup>6</sup>,  
Steven J. Ghan<sup>7</sup>, David Gregory<sup>8</sup>, Sam F. Iacobellis<sup>9</sup>, Steven K. Krueger<sup>10</sup>,  
Ulrike Lohmann<sup>11</sup>, Jon C. Petch<sup>12</sup>, David A. Randall<sup>6</sup>, Leon D. Rotstajn<sup>13</sup>,  
Richard C. J. Somerville<sup>9</sup>, Yogesh C. Sud<sup>14</sup>, Knut von Salzen<sup>15</sup>, Gregory K. Walker<sup>14</sup>,  
Audrey Wolf<sup>4</sup>, J. John Yio<sup>1</sup>, Guang Jun Zhang<sup>9</sup> and Minghua Zhang<sup>16</sup>

<sup>1</sup>Lawrence Livermore National Laboratory, USA

<sup>2</sup>NASA Langley Research Center, USA

<sup>3</sup>Observatoire Midi-Pyrenees, France

<sup>4</sup>NASA Goddard Institute for Space Studies, USA

<sup>5</sup>NOAA Geophysical Fluid Dynamics Laboratory, USA

<sup>6</sup>Colorado State University, USA

<sup>7</sup>Pacific Northwest National Laboratory, USA

<sup>8</sup>ECMWF, UK

<sup>9</sup>Scripps Institution of Oceanography, University of California, USA

<sup>10</sup>University of Utah, USA

<sup>11</sup>Dalhousie University, Canada

<sup>12</sup>Met Office, UK

<sup>13</sup>CSIRO Atmospheric Research, Australia

<sup>14</sup>Goddard Space Flight Center, USA

<sup>15</sup>Canadian Center for Climate Modeling and Analysis, University of Victoria, Canada

<sup>16</sup>State University of New York at Stony Brook, USA

Submitted to: *Q. J. R. Meteorol. Soc.*, May 8, 2001

Revised, January 25, 2002

\* Corresponding author: Atmospheric Sciences Division (L-103), Lawrence Livermore National Laboratory, P. O. Box 808, Livermore, CA 94551, USA. E-mail: [xie2@llnl.gov](mailto:xie2@llnl.gov)

## SUMMARY

This study reports the Single-Column Model (SCM) part of the Atmospheric Radiation Measurement (ARM) / the Global Energy and Water-cycle Experiment (GEWEX) Cloud System Study (GCSS) joint SCM and Cloud-Resolving Model (CRM) Case 3 intercomparison study, with a focus on evaluation of cumulus parameterizations used in SCMs. Fifteen SCMs are evaluated under summertime midlatitude continental conditions using data collected at the ARM Southern Great Plains (SGP) site during Summer 1997 Intensive Operational Period (IOP). Results from ten CRMs are also used to diagnose problems in the SCMs.

It is shown that most SCMs can generally capture well the convective events that were well developed within the SCM domain while most of them have difficulties of simulating the occurrence of those convective events that occurred only within a small part of the domain. All models significantly underestimate the surface stratiform precipitation. A third of them produce large errors in surface precipitation and thermodynamic structures. Deficiencies in convective triggering mechanisms are thought to be one of the major reasons. Using a triggering mechanism that is based on the vertical integral of parcel buoyant energy without additional appropriate constraints results in overactive convection, which in turn leads to large systematic warm/dry biases in the troposphere. It is also shown that a non-penetrative convection scheme can underestimate the depth of instability for midlatitude convection, which leads to large systematic cold/moist biases in the troposphere.

SCMs agree quantitatively well with CRMs in the updraft mass fluxes while most models significantly underestimate the downdraft mass fluxes. Neglect of mesoscale updraft and downdraft mass fluxes in the SCMs contributes to a considerable amount of the discrepancies between the SCMs and the CRMs. In addition, uncertainties in the diagnosed mass fluxes in the CRMs and deficiencies with cumulus parameterizations are not negligible.

Similar results are obtained in the sensitivity tests when different forcing approaches are used. Finally, sensitivity tests from an SCM indicate that its simulations can be greatly improved when its triggering mechanism and closure assumption are improved.

**KEYWORDS:** Cumulus Parameterization, Single Column Models, Continental Cumulus Convection

## 1. Introduction

Parameterization of physical processes associated with cumulus convection, radiation and cloud formation in general circulation models (GCMs) has proven to be extremely challenging, largely because it involves the complicated scale interactions among different processes that are not well understood. Given the large uncertainties in parameterizations, a systematic evaluation and intercomparison of the parameterization schemes used in different GCMs are an essential step in further developing and improving parameterizations. For this purpose, the ARM<sup>1</sup> Cloud Parameterization and Modeling (CPM) Working Group (WG) and the GCSS<sup>2</sup> WG 4 have conducted a series of model intercomparison case studies using data collected from the ARM Southern Great Plains (SGP) Cloud and Radiation Testbed (CART) site and other field measurements (e.g., TOGA COARE<sup>3</sup>; Webster and Lukas 1992).

The single-column model (SCM) and cloud-resolving model (CRM) are two approaches used to test and develop parameterizations in the intercomparison studies (Browning et al. 1994; Randall et al. 1996). An SCM represents a grid column of a GCM, where the large-scale advective tendencies are specified from observations and/or large-scale models and the model parameterizations are integrated forward in time. The simulated results can then be compared to observations. A CRM is a model with sufficient resolution to resolve the dynamical structures of cloud systems. It is designed to explicitly simulate the cloud-scale processes that must be parameterized in a GCM or an SCM. Since CRMs can compute some fields that are very difficult to observe, such as the vertical distributions of cumulus mass fluxes and liquid water and ice, results from CRMs are useful to diagnose problems with SCMs. However, CRMs contain parameterizations such as cloud microphysics and turbulence. Thus, their results need to be used cautiously.

The earlier intercomparison case studies include a midlatitude continental case (Case 1) conducted by ARM CPM WG and two tropical oceanic cases conducted by GCSS WG 4. The midlatitude continental case study uses data collected from the ARM SGP CART site during Summer 1995 Intensive Operational Period (IOP; Ghan et al. 2000). Ghan et al. (2000) focuses on evaluating different methodologies such as the objective analysis methods for deriving large-

---

<sup>1</sup> Atmospheric Radiation Measurement program (Stokes and Schwartz 1994).

<sup>2</sup> Global Energy and Water-cycle Experiment (GEWEX) Cloud System Study.

<sup>3</sup> Tropical Ocean Global Atmosphere -- Coupled Ocean - Atmosphere Response Experiment.

scale advective tendencies, large-scale forcing methods for use in SCMs and CRMs and the methods used to estimate surface flux forcing.

The two GCSS case studies are both based on data collected during TOGA COARE IOP. The first case study focuses on the detailed study of a squall line on a time-scale of hours. Results from CRMs are compared with kinematic and radar reflectivity observations from airborne Doppler radar (Redelsperger et al. 2000). The SCMs are forced by temperature and moisture tendencies computed from a reference CRM simulation and SCM results are evaluated by outputs from the reference CRM (Bechtold et al. 2000). The second GCSS case simulates the multi-day evolution of cloud systems during the 20-26 December 1992 subperiod, driven by the observed large-scale forcing and sea surface temperature (Krueger et al. 2002). The consensus of both SCM and CRM simulated results is compared against available observations.

In line with these earlier case studies, the ARM CPM WG and the GCSS WG 4 have conducted a joint SCM and CRM Case 3 intercomparison study, based on the data collected from the ARM SGP site during summer 1997 IOP (Cederwall et al. 2000). Compared with the earlier cases, the Case 3 study is aimed at advancing the understanding of midlatitude continental convection. It also involves more CRMs (ten) and SCMs (fifteen) compared to the earlier studies. In addition, this study breaks the IOP into several short subperiods with each lasting for 4 or 5 days, based on different weather conditions, to avoid the problem of simulations that drift away from observations over long time integration (Ghan et al. 2000) and to make the evaluation more meaningful, especially for SCMs.

The present work reports the SCM part of the Case 3 intercomparison study. The CRM results are described in a companion paper (Xu et al. 2002). The focus of the current paper is on evaluation of cumulus parameterizations used in SCMs. This is motivated by the critical role of cumulus convection in large-scale circulations. Besides, a detailed analysis of cumulus parameterizations can help better understand SCM simulations that are usually difficult to interpret because of uncertainties in both the observed large-scale input data and model parameterizations (Ghan et al. 2000).

Evaluation of individual cumulus parameterizations can be found in many early studies (e.g., Lord and Arakawa 1982; Grell et al. 1991, Xu and Arakawa 1992). Most of them use data

obtained from tropical oceanic regions, such as the GATE<sup>4</sup> dataset (Thompson et al. 1979). This is partly due to inadequate field measurements in the midlatitudes. There are significant differences between the environments of the tropical and midlatitude continental convective systems such as the underlying surface, the moisture content in the atmospheric column, the depth of the subcloud layers and instability/inhibition as summarized in Xu and Randall (2000). Extratropical summertime convection is significantly different from tropical convection. For example, extratropical convection can be formed by the lifting of a potentially unstable layer of upper air rather than of the near-surface air. Ogura and Jiang (1985) and Wu (1993) showed that deep clouds dominate the cloud population in the midlatitude convective systems. Therefore, some assumptions made in the parameterizations based on tropical observations may not be suitable for use over midlatitude continent. For example, Xie and Zhang (2000) evaluated the deep convection scheme used in the NCAR<sup>5</sup> Community Climate Model version 3 (CCM3) using the ARM SGP data. They found that the use of a positive CAPE (convective available potential energy) to trigger convection resulted in an overestimation of convection during the daytime and underestimation during the nighttime due to the strong diurnal heating that controls the variation of CAPE. Thus, validation of cumulus parameterizations against the midlatitude data is a necessary step for further improvements of cumulus parameterizations.

In this paper, we compare and evaluate cumulus parameterizations in 15 SCMs using the ARM SGP summer 1997 IOP data, which covers several continental convective events. The purpose of this paper is to systematically compare and evaluate the performance of these cumulus parameterizations under summertime midlatitude continental conditions. Through this study, we hope to identify strengths and weaknesses of these cumulus parameterizations and eventually lead to further improvement by individual participating groups.

## 2. Model descriptions

Table 1 lists the fifteen SCMs participating in the intercomparison study. Most of the SCMs are the same models that participated in the earlier case study (Ghan et al. 2000) except for CCCma1\_SP, ECMWF, GISS, MesoNH, MOUM and CCM3/SIO.

---

<sup>4</sup> Global Atmospheric Research Program (GARP) Atlantic Tropical Experiment.

<sup>5</sup> National Center for Atmospheric Research.

The parameterization schemes for deep and shallow/midlevel convection in these SCMs are listed in Table 2. It is seen that deep convection schemes used in some models are very similar. For example, the schemes used in CCCma, CCCma1\_SP, CCM3, CCM3/SIO, CCM3/SUNY and Scripps are originally proposed by Zhang and McFarlane (1995) (ZM). GFDL and McRAS use a relaxed Arakawa and Schubert scheme (Arakawa and Schubert 1974, AS) that was originally proposed by Moorthi and Suarez (1992). And the bulk mass flux scheme proposed by Gregory and Rowntree (1990) is used in CSIRO and MOUM with some modifications.

All of the deep convection schemes are based on the mass flux approach, which uses either spectral cloud ensemble models similar to AS (e.g., CSU, GFDL and McRAS) or a bulk cloud ensemble model (e.g., CSIRO, ECMWF, GISS, MesoNH, PNNL/CCM2 and MOUM). In the bulk mass flux method, only one single cloud model is used to represent an average over all cloud types within a convective ensemble. Therefore, the bulk method does not have explicit assumptions on the mass and thermodynamic budgets of sub-ensembles of cumulus clouds and does not give explicit information on the mass spectrum of various cloud types as the spectral method does. In addition, the entrainment and detrainment rates in the bulk method are often set to correctly estimate the maximum cloud-top height while they are dependent on the spectral cloud distribution in the spectral method. Yanai et al. (1976) showed that these two types of methods gave nearly identical total vertical cloud mass flux for tropical convection.

It should be noted that the ZM scheme is different from the conventional bulk mass flux schemes. It is based on the same spectral rising plume concept as used in AS. The bulk entrainment rate as a function of height is the same as a spectral model would produce because the same spectral concept is used to estimate it. However, by assuming a constant spectral distribution in cloud-base mass flux, the thermodynamic equations are reduced to the bulk form. The ZM scheme is designed primarily to treat deep convection. Therefore, some models use an additional scheme for shallow and mid-level convection. In contrast, the conventional bulk mass flux schemes are designed to represent shallow, mid-level and deep convection as function of the cloud depth and the starting level.

Other important features of these cumulus parameterizations are summarized in Table 3, such as closure assumptions, triggering mechanisms and convective downdrafts. Most models use a closure based on the assumption that CAPE is consumed by cumulus convection over a given time scale (CAPE closure). The time scale is usually set to be a few hours. CCM3/SIO revises

Table 1. Summary of SCMs used in the intercomparison study.

Model (SCM)	Model Full Name	Reference(s)
CCCma	Canadian Center for Climate Modeling and Analysis, Version 0	Lohmann et al. (1999)
CCCma1_SP	Canadian Center for Climate Modeling and Analysis, Version 1	McFarlane et al. (1992); von Salzen and McFarlane (2002)
CCM3	NCAR Community Climate Model, Version 3	Hack et al. (1998); Kiehl et al. (1996)
CCM3/SIO	Scripps Institution of Oceanography Version of CCM3	Hack et al. (1998); Kiehl et al. (1996); Zhang (2002)
CCM3/SUNY	State University of New York Version of CCM3	Hack et al. (1998); Kiehl et al. (1996); Xie and Zhang (2000)
CSU	Colorado State University	Randall and Cripe (1999)
ECMWF	European Center for Medium-range Weather Forecasts	Gregory et al. (2000) and Gregory (2001)
GFDL	NOAA Geophysical Fluid Dynamics Laboratory	See Ghan et al. (2000) for description of the GFDL model
GISS	NASA Goddard Institute for Space Studies	Del Genio and Yao (1993), and Del Genio et al. (1996)
McRAS	Microphysics of cloud/Relaxed Arakawa-Schubert scheme	Takacs et al. (1994); Sud and Walker (1999a, b)
MesoNH	Modele mesoechelle non-hydrostatique	Lafore et al. (2000)
PNNL/CCM2	Pacific Northwest National Laboratory Version of CCM2	Hack et al. (1993); Ghan et al. (1997)
Scripps	Scripps Institution of Oceanography	Iacobellis and Somerville (1999)
MOUM	Met Office Unified Model	Pope et al. (2000)

Table 2. Summary of deep and shallow/midlevel convection schemes used in the SCMs.

Model	Deep Convection Scheme	Shallow /midlevel Convection Scheme
CCCma	Revised ZM (Lohmann, personal communication)	No
CCCma1_SP	Prognostic ZM (von Salzen, personal communication)	von Salzen and McFarlane (2002)
CCM3	ZM (Zhang and McFarlane 1995)	Hack (1994)
CCM3/SIO	Revised ZM (Zhang 2002)	Same as CCM3
CCM3/SUNY	Revised ZM (Xie and Zhang 2000)	Same as CCM3
CSIRO	Bulk mass Flux (Gregory and Rowntree 1990; Gregory 1995)	The deep convection scheme applicable to all types of convection
CSU	Prognostic AS (Ding and Randall 1998)	No
ECMWF	Bulk mass flux (Tiedtke 1989; Gregory et al. 2000; Gregory and Guichard, 2001)	Mass flux treatment but different closure/specification compared to deep convection
GFDL	Relaxed AS (Moorthi and Suarez 1992)	No
GISS	Bulk mass flux (DelGenio and Yao 1993)	The deep convection scheme applicable to all types of convection
McRAS	Revised relaxed AS (Sud and Walker 1993)	No
MesoNH	Bulk mass flux (Bechtold et al. 2001)	The deep convection scheme applicable to all types of convection
PNNL/CCM2	Bulk mass flux (Hack 1994)	The deep convection scheme applicable to all types of convection
Scripps	ZM	No
MOUM	Bulk mass flux (Gregory and Rowntree 1990; Gregory 1995)	The deep convection scheme applicable to all types of convection



Table 3. Summary of some relevant aspects of the deep convection schemes. CWF is cloud work function defined by Arakawa and Schubert (1974), while  $CWF_{clim}$  is the climatological value of CWF. CAPE is convective available potential energy. LCL is the lifting condensation level.

Model	Closure	Triggering mechanism	Downdrafts
CCCma	CAPE	$CAPE > 0$ and Buoyancy at $LCL > 0.1K$	Convective precipitation induced downdrafts. Downdrafts starting at or below the updraft detrainment layer. No downdrafts below cloud base
CCCma1_SP	Prognostic closure	$CAPE > 0$	Same as CCCma
CCM3	CAPE	$CAPE > 0$	Same as CCCma
CCM3/SIO	CAPE generation rate	CAPE generation rate $> 0$	Same as CCCma
CCM3/SUNY	CAPE	Positive dynamic CAPE	Same as CCCma
CSIRO	A stability-dependent mass flux	Parcel buoyancy	Downdrafts exist if the updraft is saturated and the depth of cloud layer exceeds 150 hPa. Downdrafts not allowed within 100 hPa of the surface
CSU	A prognostic closure using the cumulus kinetic energy	$CWF > 0$ and constraints on the entrainment rate and the saturation of cloudy air at the cloud top	No
ECMWF	CAPE	Parcel buoyancy	Convective precipitation induced downdrafts.
GFDL	Same as McRAS	$CWF > 1.4 * CWF_{clim}$ for the deepest cloud type, $CWF > CWF_{clim}$ for other cloud types	No
GISS	Cloud base neutral buoyancy	Parcel buoyancy	Downdrafts specified to be a third of updrafts
McRAS	Relax the state toward quasi-equilibrium	$CWF > 0$ and $RH > RH_{crit}$	Rain-evaporation induced downdrafts
MesoNH	CAPE	Parcel buoyancy and vertical motion	Convective precipitation induced downdrafts
PNNL/CCM2	A stability-dependent mass flux	Parcel buoyancy	No
Scripps	CAPE	$CAPE > 0$	Same as CCCma
MOUM	CAPE	Parcel buoyancy	Similar to CSIRO

the CAPE closure for ZM scheme so that the cloud-base mass flux is determined by the CAPE generation rate due to the large-scale tropospheric forcing (Zhang 2002). CSU employs a prognostic closure using the cumulus kinetic energy (Pan and Randall 1998) for AS scheme. CCCma1\_SP uses a similar prognostic closure suggested by Pan and Randall (1998) to account for the effects of self-organization of convective systems for ZM scheme (von Salzen, personal communication). Some of the bulk mass flux schemes use a closure based on a stability-dependent mass flux (i.e., CSIRO, GISS and PNNL/CCM2).

The primary triggering mechanisms used in these cumulus parameterizations are either based on CAPE (or CWF<sup>6</sup>), or based on local parcel buoyancy. The former measures the instability based on the vertical integral of parcel buoyant energy while the latter measures the instability by lifting up a parcel to a specific distance or between two model layers. Most schemes with the CAPE/CWF triggers need additional constraints as shown in Table 3. For example, CCM3/SUNY requires a positive contribution from large-scale dynamic processes to the existing positive CAPE (Xie and Zhang 2000). McRAS uses the relative humidity (RH) that exceeds 0.9 times the critical value  $RH_{crit}$  for stratiform condensation, in addition to the positive CWF.

Another feature seen from Table 3 is that convective-scale precipitation-driven downdrafts are parameterized in most of the models except for PNNL/CCM2, CSU and GFDL. In addition, almost all models treat convection as penetrative convection except for the Hack scheme (Hack 1994) used in PNNL/CCM2. It uses a three-level non-entraining cloud model and convection couples together triplets of layers.

There are also many differences among the SCMs in parameterizing physical processes such as stratiform cloud formation, radiation and turbulent processes. For example, most models use a prognostic scheme to predict stratiform cloud condensate, except for CCCma1\_SP, CCM3, CCM3/SIO and CCM3/SUNY, which use a diagnostic scheme to estimate cloud fraction and cloud condensate. Some models (CSU, ECMWF, GFDL, GISS, McRAS, MesoNH, PNNL/CCM2 and Scripps) consider the detrainment of condensate water from cumulus clouds in their prognostic cloud scheme. Radiation schemes in most models are from either the ECMWF<sup>7</sup> model or the NCAR CCM. They employ a variety of turbulent parameterization schemes. Detailed descriptions of these schemes can be found in the references listed in Table 1.

---

<sup>6</sup> Cloud work function (CWF) as defined by Arakawa and Schubert (1974) is a function of cloud types and it is equal to CAPE when the entrainment rate is zero.

### 3. Data

The data used to drive and evaluate the SCMs and CRMs were collected at the ARM SGP site during summer 1997 IOP, from 18 June (2330 UTC, Julian day 170) to 17 July (2330 UTC, Julian Day 199, hereafter, Day is used instead of Julian day). Observations at the ARM site used to characterize the atmospheric column include 3-hourly soundings at 5 locations, surface meteorology network, wind profilers data, satellite data and radar rainfall estimates; details are given in Ghan et al. (2000). The horizontal and vertical advective tendencies of temperature and moisture and vertical velocity are derived from the constrained variational analysis (Zhang and Lin 1997; Zhang et al. 2001). In the analysis, the atmospheric state variables are forced to satisfy the conservation of mass, heat, moisture and momentum. Ghan et al. (2000) showed that the variational analysis approach significantly improved the quality of the input data, and thereby the SCM simulations.

The summer 1997 IOP contained a wide range of summertime weather conditions. Associated with the activities of the large-scale upper-level troughs and ridges over the North America continent, the ARM SGP site experienced several intensive precipitation events and dry and clear days during this IOP (Fig. 1). The three subperiods defined by A, B and C in Fig. 1, with each including two or three precipitation events and lasting for 4 or 5 days, are selected to test model performance. To help illustrate the synoptic conditions during the three subperiods, Figure 2 shows a series of 4 km resolution infrared cloud images, corresponding to the peaks of the eight strong precipitation events that occurred on Days 179, 180-181, 190, 191, 192, 195, 196 and 197-198, from the seventh NOAA GOES<sup>8</sup> (GOES-7) measurements. The circle in Fig. 2 is the variational analysis domain or the SCM domain.

Subcase A features a weak precipitation event occurred on Day 179 and a strong precipitation event on Days 180-181. The weak precipitation event was associated with very moist air along with the tail end of a front moving eastward across the SGP site. Several localized individual thunderstorms developed in early morning in the southern part of the SCM domain. However, the domain-averaged precipitation was very weak. The heavy precipitation event on Days 180-181 was mainly associated with a complex of thunderstorms that developed

---

<sup>7</sup> European Centre for Medium-range Weather Forecasts.

<sup>8</sup> National Oceanic and Atmospheric Administration Geostationary Operational Environmental Satellite.

in southeastern Kansas in the late evening of Day 180 (local time) and moved southeastward across the SCM domain during Days 180-181. Here local noon corresponds to 1700 UTC. The strong convective complex affected the major parts of the SCM domain. This convective event was well captured by the ARM sounding array.

Subcase B contained three precipitation events. The first one (Day 190) was related to a complex of thunderstorms that developed in southeastern Kansas during the evening hours of Day 190 and moved southeastward into southwestern Missouri and northeastern Oklahoma overnight. Only the northern edge of the ARM sounding array experienced heavy rainfall based on the Arkansan Basin Red River Forecast Center (ABRFC) 4 km WSR-88D radar precipitation estimates (not shown). The other two precipitation events (Days 191 and 192) were more localized. They were associated with a developing upper-level trough that extended southeastward over the North America continent. The southern edge of the trough moved across the SCM domain and brought in a moist and unstable air mass to the area. The large-scale circulation and diurnal heating allow isolated-scattered showers and thunderstorms to develop within the SCM domain in late afternoon and evening during Days 191 and 192.

Subcase C experienced one very weak precipitation event on Day 195 and two consecutive precipitation events over the last three days. The weak precipitation event was associated with insolated thunderstorms along with a slowly moving cold front. The convective system developed within the SCM domain, however, it only occupied a small part of the SCM domain. Most of the SCM domain was clear. The second precipitation event on Day 196 was associated with a trough of low pressure moving eastward into the SGP site. A complex of thunderstorms started to develop early in the morning of Day 196 at the northwestern part of the SCM domain and then slowly moved over the entire domain. The third precipitation event was associated with a nearly stationary upper-level trough that developed and extended southeastward across the SGP site. Thunderstorms developed during the evening hours between Wichita, Kansas and Ponca City, Oklahoma and continued to rain over the same general area for about 12 hours. Northeastern part of the domain experienced heavy rainfall.

As described above, for the eight precipitation events of the three subcases, some of the convective events were well located within the SCM domain and some were not. For those cases that only a small portion of the SCM domain experienced heavy rainfall, the SGP sounding array may not properly capture the characteristics of the strong convective complex.

## 4. Intercomparison and interpretation of simulation results from the baseline experiment

The baseline experiment in this study uses the horizontal and vertical advective tendencies of temperature and moisture and surface fluxes specified from the observations and the radiative heating rates are calculated from model parameterizations. See Cederwall et al. (2000) for the detailed description of experiment design. Figure 3 shows the total large-scale advective tendencies of temperature and moisture and the corresponding surface precipitation rates for Subcases A, B and C. It is seen that the forcing is very strong during the strong precipitation event occurred on Day 181 in Subcase A and it is relatively weak during other precipitation periods. In general, the observed precipitation events correspond to the large-scale advective cooling in the middle and upper troposphere (maximum centered at 400 – 450 hPa) and moistening in the lower troposphere (maximum centered at about 750 hPa), however, for the first precipitation events in Subcases B and C, the upper level advective cooling is actually associated with weak low-level moisture divergence. Recall that only small portions of the SCM domain experienced precipitation during these two precipitation events. This suggests that the domain averaged large-scale forcing may not be able to properly capture the typical thermodynamic structure of a strong convective complex.

### a. Surface precipitation rates

Surface precipitation rates are closely related to model cumulus convection schemes and therefore they are evaluated first. The satellite images clearly show that the precipitation events during these subperiods were associated with mesoscale convective systems that were influenced by the large-scale circulations. To quantitatively demonstrate the relative importance of cumulus clouds and stratiform clouds in these convective events, we partitioned the total rainfall into convective and stratiform components based on the ABRFC 4 km radar data using the algorithm developed by Johnson and Hamilton (1988). They used  $6 \text{ mm hr}^{-1}$  as the threshold for convective-stratiform partitioning of an OK PRE-STORM<sup>9</sup> squall line.

Most of the convective events are dominated by cumulus precipitation, especially for the strong precipitation event that occurs on Days 180-181 where more than 80% of surface rainfalls

---

<sup>9</sup> Oklahoma-Kansas Preliminary Regional Experiment for STormscale Operational and Research Meteorology.

are from cumulus clouds (Table 4). The exceptions are the first precipitation event in Subcase A and the third precipitation event in Subcase B, in which cumulus clouds and stratiform clouds have comparable contributions to the total precipitation. For all three subperiods combined, 72% of surface rainfalls are from cumulus clouds, but 28% are from stratiform clouds.

Table 4. Time mean surface precipitation rates ( $\text{mm day}^{-1}$ ). Letters A, B and C in parenthesis denote Subcases A, B and C, respectively.

Convective Events	Total	Cumulus	Stratiform
Day 179 (A)	4.14	2.17	1.97
Days 180-181 (A)	24.7	20.1	4.6
Day 190 (B)	9.33	7.20	2.13
Days 191 (B)	10.54	7.13	3.41
Days 192 (B)	7.27	3.61	3.76
Day 195 (C)	1.96	1.56	0.40
Days 196-197 (C)	6.30	4.97	1.33
Days 197-198 (C)	10.0	6.92	3.07
Total mean	9.28	6.70	2.58

Figures 4a-f show time series of the total and stratiform precipitation rates estimated from the radar data (black line) and simulated from 13 SCMs for Subcases A, B and C, respectively. Note that results from CCM3/SIO and CCM3/SUNY are not shown in the baseline experiment. Because these two models are identical to CCM3 except for some details of cumulus parameterizations, their results will be shown in section 5 to illustrate the sensitivity of CCM3 simulations to different convective triggering mechanisms and closure assumptions.

It is seen that most models simulate reasonably well the strong precipitation event (Days 180-181) in Subcase A and the last two precipitation events (Days 191-192) in Subcase B although they are less successful in simulating the rest of convective events (Figs. 4a-c). A noticeable feature appearing in Figs. 4a-c is that some models show large spurious precipitation during non-precipitation periods. This is especially true for the models that use positive CAPE only as a triggering mechanism (see Table 3), such as CCM3, Scripps and CCCma1\_SP. They produce pronounced diurnal variations of surface precipitation that are not observed. This is less serious in those models with triggering mechanisms based on CAPE/CWF with additional

constraints or the parcel buoyancy methods, except for CSU, in which convection is occasionally too active. It is also noticed that CCCma largely underestimates most of the precipitation events. This suggests that the revised triggering mechanism for ZM used in CCCma overly inhibits convection. PNNL/CCM2 also underestimates the magnitudes of most of the precipitation events except for the last two events in Subcase B.

Figures 4d-f show that stratiform precipitation rates are largely underestimated in all SCMs. For most of the convective events, there is little stratiform precipitation produced in most models. Only ECMWF and GISS produce reasonable stratiform precipitation amounts during the strong precipitation event in Subcase A. This is partially because cumulus convection is too active in most models. Cumulus convection consumes most of the moisture transported from the boundary layer. Assumptions on detrainment of condensate water from cumulus clouds could also contribute to this problem.

Table 5 shows the time-averaged total precipitation rates for Subcases A, B and C. Consistent with the discussion above, CCM3, Scripps, CCCma1\_SP, MesoNH (Subcase B only) and CSU (Subcase B only) greatly overestimate the observed surface precipitation rates while CCCma, ECMWF, MesoNH (Subcase A only) and PNNL/CCM2 greatly underestimate the observed values of all subperiods. The rest of the models have the time-averaged precipitation rates within  $1 \text{ mm day}^{-1}$  of the observations.

To quantify the statistical similarity among the models, Figs. 5a-b present a Taylor diagram (Taylor 2001) for the total and convective surface precipitation fields, respectively. In Fig. 5, points A1 to A3 represent the models that use CAPE only triggers, points B1 to B4 represent the models that use CAPE/CWF triggers with additional constraints, points C1 to C5 represent the models that use the parcel buoyancy triggers, point X represents the consensus of ten CRMs results (Xu et al. 2002), and point Obs represents the observations. Note that CRM consensus for convective precipitation is not shown because only a few models provide the data. For simplicity, here we only show results for Subcase B. Results from Subcase A (not shown) for most models except CCCma show much better agreement with the observations than those from Subcase B. This is because Subcase A is characterized by a strong precipitation event associated with a strong large-scale forcing (Figs. 3a, d and g) and most schemes could well simulate the convection in such environments. Results from Subcase C (not shown) are similar to Subcase B.

Table 5 Time mean precipitation rates during Subperiods A, B and C.

Models	Subcase A	Subcase B	Subcase C
Observation	8.21	4.17	4.72
CCM3	13.18	10.88	9.28
Scripps	10.83	6.98	7.09
CCCma	3.25	1.62	1.69
CCCma1_SP	11.76	5.93	6.92
CSU	8.78	6.05	5.56
McRAS	9.10	4.76	4.45
GFDL	8.79	4.92	4.38
ECMWF	6.88	N/A	3.05
GISS	8.37	5.07	4.56
CSIRO	8.41	4.91	4.00
MOUM	8.92	5.00	5.54
PNNL/CCM2	5.63	3.23	2.86
MesoNH	6.98	6.1	4.71

Comparing Fig. 5a with 5b, almost all models except CCCma (B1) show better agreement with the observations for the total precipitation field than its convective part, in terms of the RMS error and the amplitude of temporal variability, while the correlation coefficients are similar for both fields. Another feature is that most models except CCCma overestimate the observed temporal variations in the convective precipitation field. This is consistent with the fact that most models overestimate the convective precipitation and underestimate the stratiform precipitation discussed earlier.

In general, the models that use the CAPE only triggers and CSU (point B2) produce the least agreement with the observations. The rest of the models generally produce comparable results to the CRM consensus (point X). Note that the CRMs have much smaller intermodel differences in the surface precipitation field than those seen in the SCMs and the first precipitation event is severely delayed in Subcase B (Xu et al. 2002). The comparable results suggest that these SCMs are doing as well as can be expected given uncertainties in the forcing and the observed rainfall.



It is also noteworthy that the correlation coefficients in these models except GFDL and PNNL/CCM2 are also quite small (less than 0.6). The small coefficients are partially related to the failure of promptly initiating the first convective event of Subcase B by these SCMs (Fig. 4b). This is also the case in all CRM simulations (Xu et al. 2002). The delayed convection may be caused by the dry thermodynamic profiles at the beginning of the event as discussed below.

SCMs/CRMs are driven by the SCM domain-averaged large-scale forcing, however, the forcing may not be able to capture well those convective events that only occurred in a fraction of the SCM domain, such as the first precipitation events of the three subcases (Fig. 2). Figures 6a-d respectively show the vertical profiles of moist static energy (MSE, solid line) and saturated MSE (dashed line) at the beginnings of four selected precipitation events occurred on Days 180-181, 190, 191 and 195 of Subcases A, B and C, with the sounding profiles at the station where convection is occurring. Figs. 6e-h are the same as Figs. 6a-d except they are calculated from the SCM domain average profiles. The MSE values are normalized by  $C_p$  ( $1004 \text{ J kg}^{-1} \text{ K}^{-1}$ ), the specific heat at constant pressure. Comparing Figs. 6a-d with Figs. 6e-h, it is much easier to diagnose convection from the single-sounding profiles than from the SCM domain-averaged profiles. Another noticeable feature is that sometimes convection is triggered at higher levels (e.g., probably Day 191, Figs. 6c, g) and some models may not handle that well.

Figure 6 also helps to understand why most models perform well for some convective events and perform worse for the others. We have shown in Fig. 4 that most models typically performed well for the convective events that occurred on Days 180-181 in Subcase A and Days 191-192 in Subcase B. Recall that these convective events were well developed within the ARM SCM domain and most part of the domain experienced rainfall. Thus, the mean thermodynamic profiles can properly describe the characteristics of convective systems. As shown in Figs. 6e, g, which correspond to these convective events, the domain averaged thermodynamic profiles show that the environmental atmospheres are conditionally unstable and there is enough moisture at low levels to make a parcel lifted a reasonable distance buoyant. In contrast, for the convective cases in which only a part of the domain was affected by convective systems, such as the first precipitation events of Subcases B and C, the average profiles (Figs. 6f, h) are so dry at the low levels that few cumulus parameterizations would be triggered. This indicates that the problem with SCMs that are trying to validate against the timing of specific convective events is that without knowledge of the probability distribution functions (pdfs) of temperature and moisture in

the planetary boundary layer (PBL), it is impossible to tell that a profile that is not overly conducive to convection in the mean might not have a tail of the distribution within the gridbox that is conducive. It suggests a need to use CRMs to develop pdfs of PBL properties that might give SCMs insights into how to get the statistics of convective occurrence right.

The aforementioned discussions on the timing of simulated precipitation also suggest that deficiencies in convective triggering mechanisms are responsible for the precipitation errors in some models, such as CCM3, Scripps, CCCMa1\_SP, CSU and CCCma. It is also noticed that the triggering mechanisms used in these models are all primarily based on CAPE/CWF. The surface sensible and latent heat fluxes can largely affect the generation of CAPE over the land surface where these fluxes are large and have strong diurnal variations. Over these areas, convection could be more active for schemes using CAPE/CWF than those using parcel buoyancy as triggering mechanisms if no any additional constraints are used.

To help discuss this issue, we analyze CAPE and CIN (convective inhibition) based on the ARM observations for Subcases B and C (Fig. 7). The observed surface precipitation rates are also shown. CAPE is calculated under the assumption that an air parcel ascends along a reversible moist adiabat with the originating level at the surface. CIN is the negative value of CAPE value between the free convection level and the surface.

It is seen that CAPE exhibits strong diurnal variation with maximum during daytime and minimum during nighttime due to the strong solar diurnal cycle over the land surface. The opposite is true for CIN that is usually large during non-precipitating periods. A large CIN value prevents convection from consuming the large amount of CAPE existed during these periods. Therefore, a positive CAPE is not a sufficient condition to trigger convection. In fact, there are obvious phase shifts between the observed CAPE and surface precipitation rates since most of the convective events occur in late evening and early morning. Therefore, it is not surprising to see the excessive initiation of convection in those models that use the CAPE triggers only to diagnose the onset of convection. It should be noted that the triggering problem over tropical oceans might not be as serious as over the midlatitude land because the diurnal variations of CAPE is weaker due to the large heat capacity of ocean.

## **b. Cloud fractions and outgoing longwave radiation fluxes**

Because all SCMs include cloud parameterizations, in addition to cumulus parameterizations, it is necessary to show how the clouds and radiation are impacted by some deficiencies in cumulus parameterizations discussed earlier. Similar to Fig. 5, Fig. 8a presents a Taylor diagram for comparing the statistical similarities and differences of simulated column cloud fractions among the SCMs for Subcase B. Note that cloud fraction data is not available for CSU (point B2). The observed cloud fraction is from the retrievals of ground-based millimeter wave cloud radar (MMCR; Moran et al. 1998) at the ARM central facility. Time series of the observations is shown in Xu et al. (2002), which indicates rather large temporal variations for this subcase.

Most SCMs show large variabilities as in the observations except for CCM3 (point A1) and CCCma1\_SP (point A3), which produce nearly overcast conditions at most times. As mentioned earlier, no prognostic cloud parameterization is implemented in these two models. Another noticeable feature shown in Fig. 8a is that there are large differences among the models even they use similar cumulus parameterizations. This implies that the details in the cloud parameterizations impact the simulations of cloud fraction. Comparing to the consensus of CRM results, the SCMs generally show better simulations in the amplitude of the temporal variability in column clouds and larger RMS errors and lower correlation with the observations.

The observed outgoing longwave radiative fluxes (OLR) at the top of the atmosphere (TOA) from GOES satellite observations are compared with simulations in Fig. 8b for Subcase B also. Note that CRM results are not available because of the prescribed radiation in their simulations. It is seen that most models exaggerate the amplitude of the temporal variability than the observed while CCM3 (point A1) and Scripps (point A2) underestimate the temporal variability. Differences in OLR among the SCMs are mainly in the amplitude of the temporal variabilities, compared to surface precipitation rates and column cloud fractions (Fig. 5 and Fig. 8a). This suggests that the impact of convection and stratiform cloud on the radiative energy is significant in the context of SCM modeling approach.

### **c. Temperature and moisture structures**

The simulated temperature and moisture biases (Fig. 9) are averaged between pressure levels 115-915 hPa to minimize their influence by PBL processes. A noticeable feature seen in Fig. 9 is that the model simulations start to diverge after a few hours of integration. The models with excessive precipitation, such as CCM3, Scripps and CSU (Subcases B and C only), generally

produce larger warm/dry biases. The excessive convective activities result in more convective heating and drying. CCCma also shows rather large warm/dry biases during most of the periods even though the model largely underestimates the observed convective events. The reason is unclear and needs to be investigated further. On the other hand, PNNL/CCM2 shows rather large cold and moist biases. This is mainly related to the non-penetrative convection scheme used in the model. As shown later, a non-penetrative convection scheme largely underestimates the depth of instability and therefore leads to large cold and moist biases. For other models, temperature and moisture biases are relatively small for the layer-mean values.

Next, the vertical structures of the temperature and moisture departures from observations, averaged over the three subperiods, are shown in Fig. 10. For temperature (Figs. 10a-c), most SCMs generally produce smaller errors in the middle troposphere and slightly larger errors in the lower and upper troposphere. The exceptions are CCM3 and Scripps, which show large warm biases throughout the troposphere. The largest intermodel difference appears in the upper troposphere, where the models with the ZM and AS types of convection schemes generally produce warmer atmosphere than those with the conventional bulk mass flux schemes (see Table 2). CCM3 and PNNL/CCM2 are the two typical cases: the former shows the largest warm biases and the latter shows the largest cold biases in the middle and upper troposphere. It is also noteworthy that a majority of models produce warm biases near the surface, suggesting possible errors in the specified surface fluxes and surface temperature. Besides, deficiencies in PBL parameterizations and weak downdrafts in SCMs (see Section 4d) may also account for the large errors near the surface.

For moisture (Figs. 10d-f), McRAS, GFDL, GISS and ECMWF have rather small biases in the entire troposphere, comparable to CRMs (Xu et al. 2002). The simulations are largely diverse below 765 hPa, generally with dry biases in those having warm biases and moist biases in those having cold biases. CCM3 and Scripps have particularly large dry biases between 765 – 865 hPa and PNNL/CCM2 has slightly larger moist biases below 765 hPa. It is also noted that CCCma has extremely large moist biases near the surface. This is related to the large moistening produced from its turbulent parameterization (Lohmann, 2001; personal communication).

#### **d. Cloud mass flux**

Cloud mass fluxes are highly related to convective heating/drying profiles; however, they are usually not easy to observe. In this study, we make use of the cloud mass fluxes diagnosed from CRMs to evaluate those produced from SCMs. The criteria for diagnosing updraft ( $M_u$ ), downdraft ( $M_d$ ) and net mass fluxes ( $M_c$ ) from CRMs data are given in Xu et al. (2002).

Figures 11a-c respectively compare the vertical structure of updraft ( $M_u$ ), downdraft ( $M_d$ ) and net ( $M_c$ ) cloud mass fluxes estimated from the SCMs with those diagnosed from the CRMs averaged over precipitation periods (observed precipitation rate  $> 0.36 \text{ mm day}^{-1}$ ) of Subcases A, B and C. Figures 11d-f are identical to Figs. 11a-c except for non-precipitation periods (observed precipitation rate  $< 0.36 \text{ mm day}^{-1}$ ). Note that the mass flux data are not available for GFDL and ECMWF. It should be also noted that the radiative heating rate is specified from the ECMWF analysis in the CRM simulations. The black solid lines in Figs. 11c, f are the observed large-scale mean mass flux ( $M = -?$ ). A detailed description of individual CRM results can be found in Xu et al. (2002). Data for the updraft and downdraft mass fluxes are not available for some SCMs.

During precipitation periods (Figs. 11a-c), the CRMs show large  $M_u$  in the middle and upper troposphere between 300 – 700 hPa. Downdraft mass flux  $M_d$  has a comparable magnitude with  $M_u$ . The maximum  $M_d$  is around 600 hPa. As a result, the net mass flux  $M_c$  is relatively small compared to the updraft and downdraft mass fluxes. It is positive in the middle and upper troposphere and negative in the lower troposphere. As shown in Raymond (1993), this is a typical vertical structure of the net cloud mass fluxes in cumulonimbus clouds in midlatitudes where cloud base heights and PBL depths are usually higher and the precipitation induced downdrafts can equal or exceed updrafts in the lower parts of cumulonimbi.

Compared to the CRMs, the updraft mass flux in the SCMs is smaller in the middle and upper troposphere and larger in the lower troposphere. The latter is associated with lower cloud base heights in SCMs. The downdrafts in the SCMs are very weak. This leads to much larger net mass fluxes in virtually all SCMs, especially in the lower troposphere. Most models fail to produce the negative mass flux shown in the CRMs due to the weak downdrafts and lower cloud base heights. Only MesoNH exhibits small negative mass flux in the lower troposphere because of its higher cloud base heights.

It should be noted that there are differences in defining cloud mass fluxes between CRMs and SCMs. In the CRMs, the cloud mass fluxes contain all types of updrafts and downdrafts

including convective-scale and mesoscale updrafts and downdrafts. In contrast, the SCMs only parameterize convective-scale updrafts and convective-scale precipitation-induced downdrafts (see Table 3). This can partly explain why the differences are so large. Many studies (e.g., Johnson 1980; Houze and Cheng 1981; Cheng and Yanni 1989) have shown that mass fluxes in mesoscale updrafts and downdrafts are significant relative to convective mass fluxes.

The mesoscale and convective scale mass fluxes diagnosed from the UCLA/CSU CRM are used to discuss the importance of mesoscale updrafts and downdrafts (Fig. 12). These data are not available from other CRMs. The mass fluxes are averaged over the same precipitation periods as those in Figs. 11a-c. The partitioning method used for this analysis is described in Xu (1995). The maximum draft velocity below the melting level is the main variable used to distinguish convective regions from stratiform regions. An entire CRM column is classified into a convective region if the maximum draft velocity is, at least, twice the averaged value of two adjacent grid columns, or exceeds  $3 \text{ m s}^{-1}$ , or surface precipitation rate exceeds  $25 \text{ mm h}^{-1}$ . Some other variables are also used to diagnose shallow convection columns. In addition, some gravity wave contributions are eliminated from the mass flux profiles by raising the thresholds on the condensate mixing ratios to  $3 \times 10^{-4} \text{ g g}^{-1}$  when draft velocities are less than  $1 \text{ m s}^{-1}$ . Table 6 provides time mean (all three subcase combined) total, convective and stratiform precipitation rates, as well as the ratio of the convective and stratiform precipitation to the total precipitation, for the CRM and ABRFC radar estimate. It is seen that the CRM produced surface precipitation rates are in an excellent agreement with the observations. The stratiform and convective components from the CRM are also very close to the observational estimates. This provides confidence on the partition of the convective-scale and mesoscale mass fluxes from this CRM.

Table 6: Surface precipitation rates ( $\text{mm day}^{-1}$ ) averaged over Subcases A, B and C

	UCLA/CSU CRM	Observations
Total	5.56	5.51
Convective	4.04	3.98
Stratiform	1.52	1.53
Convective/Total	0.73	0.72
Stratiform/Total	0.27	0.28

The UCLA/CSU CRM results indicate that convective-scale updrafts are a major contributor to the total updraft mass flux at levels below 300 hPa (Fig. 12a). Its maximum mass flux appears in the middle troposphere ( $\sim 600$  hPa) while the counterpart of mesoscale updrafts appears in the upper troposphere ( $\sim 300$  hPa). In contrast to the updrafts, mesoscale downdrafts have comparable values with convective-scale downdrafts. The convective-scale downdrafts are mainly located in the lower troposphere. For the net mass fluxes, the convective-scale mass flux is dominated by the convective-scale updrafts throughout the entire troposphere except for the levels below 800 hPa while the mesoscale mass flux is dominated by the mesoscale downdrafts in the lower and middle troposphere. It is seen from Figs. 11a-b that the SCMs generally capture well the convective-scale updrafts but most of the SCMs still significantly underestimate the convective-scale downdrafts except for GISS. It is worth noting that most SCMs have higher maximum convective-scale updraft levels (400 hPa) than the CRM (600 hPa). Since mass fluxes diagnosed from CRMs have noticeable intermodel differences as shown in Fig. 11, it is not possible to draw firm conclusions based on results from one CRM. The partitioning results from other CRMs need to be investigated.

It should be noted that large uncertainties remain in the mass flux diagnosed from the CRMs and the methodology used for partitioning the total mass flux into convective-scale and mesoscale components. The presence of these uncertainties will not allow one to fully explain the large discrepancies between the SCMs and the CRMs. For example, Xu et al. (2002) show that the diagnosed mass fluxes in the CRMs include contributions not only from convective-scale and mesoscale circulations, but also from gravity waves since many different scales of motions are present in CRM simulations. Therefore, some of the strong updrafts and downdrafts in the upper troposphere may be related to unrealistically, strong gravity wave activities in the simulations.

The weak downdrafts in SCMs are also related to the assumptions made in the parameterizations. For example, downdrafts in SCMs are limited not to exceed certain amounts of the corresponding updrafts. Some schemes (e.g., ZM) assume that no downdrafts are allowed below cloud base (see Table 3). In addition, too strong downdrafts may overly suppress subsequent convection. An independent study (Bechtold, 2001; personal communication) shows that downdrafts can be tuned to give the right strength and good lower level temperature and moisture biases corresponding to the CRM results, but they got too low precipitation.

During non-precipitation periods (Figs. 11d-f), the updraft, downdraft and net cloud mass fluxes diagnosed from all CRMs are close to zero. However, some SCMs such as CCM3, Scripps, CSU and CCCma1\_SP generate large cloud mass flux during non-precipitation periods. This is closely related to the problem with triggering of convection, as discussed earlier.

Figure 11 also shows considerable intermodel differences among the SCMs. In general, the CCM3, Scripps, CCCma1\_SP and CSU produce smaller (larger) mass fluxes than the others during precipitation periods (non-precipitation periods). This suggests that the warm and dry biases found in these models are mainly related to the spurious convections generated over the non-precipitation periods. It is noted that  $M_c$  in PNNL/CCM2 (Fig. 11c) is significantly small in the upper troposphere in comparison with other SCMs. The  $M_c$  is mainly located in the middle and lower troposphere with the maximum value around 600 hPa. Compared to other SCMs, PNNL/CCM2 typically has a higher cloud base and a lower cloud top. These imply that the model, with a non-penetrative convection scheme, underestimates the depth of instability. It is also noted that CCCma produces the weakest cloud mass flux. This is consistent with the fact that this model overly suppresses convection as shown earlier.

#### e. Q1 and Q2

The time-averaged apparent heat sources (Q1, Yanai et al. 1973) and contributions from individual physical processes such as radiative heating (Qr), cumulus convection (Q1c) and turbulent processes (Q1v), over precipitation and non-precipitation periods of the three subcases are examined next. Note that the contributions from individual physical processes are not available from some models. During precipitation periods, the observed Q1 shows strong heating in the upper troposphere and weak cooling below 765 hPa (Fig. 13a). The maximum heating is at about 400 hPa. Most models well reproduce the observed Q1 above 700 hPa while CCM3, Scripps and CCCma1\_SP overestimate the heating in the lower levels. Consistent with the weak mass fluxes, PNNL/CCM2 largely underestimates the strong heating in the upper troposphere. This contributes to its large cold biases seen in the upper troposphere.

It is also seen that convective heating (Q1c) contributes the most to the total Q1 for the levels above 865 hPa for all models. Below 865 hPa, radiative cooling (Qr) and turbulent processes (Q1v) also have significant contributions to the total Q1 (Figs. 13 b-d). Consistent with the small



stratiform precipitation produced in the models, contributions from stratiform clouds are very small (not shown).

During non-precipitation periods, the observations show very weak heating and cooling (Fig. 14a). Compared to the observations, the models with the CAPE only triggers and CSU produce excessive warming in the middle and upper troposphere. This excessive warming is a major contributor to the warm biases seen in Figs. 10a-c. The individual components clearly show that model cumulus convection (Q1c) is the major reason for the excessive warming. In fact, the intermodel differences are small in other components compared to those in Q1c except for the boundary layer where Q1v also shows rather large intermodel differences (Figs. 14b-d).

The time-averaged Q2 and the contributions from individual physical processes such as cumulus convection (Q2c) and turbulent processes (Q2v), over precipitation and non-precipitation periods of the three subcases, are shown in Figs. 15 and 16, respectively. Q2 is the apparent moisture sinks introduced by Yanai et al. (1973), representing moisture changes due to condensation, evaporation and subgrid transport. During precipitation periods, the calculated Q2 from the models agrees closely with the observed Q2 above 765 hPa while the models diverge below (Fig. 15a). In general, the models with the CAPE only triggers show excessive drying and the others show excessive moistening. It is seen that convective drying is a major contributor to the total drying above 750 hPa (Fig. 15b). Below that level, drying from Q2c and moistening from Q2v have comparable contributions but opposite sign to the total Q2 (Fig. 15c). During non-precipitation periods (Fig. 16), these models show considerable intermodel differences in Q2. The models with CAPE only triggers generally produce strong drying in the middle and lower troposphere. This is mainly related to the excessive convective drying (Q2c) generated in these models during non-precipitation periods. They are consistent with the results seen in the simulated moisture field (Figs. 10d-f).

## 5. Sensitivity tests

In the previous section, we have discussed the simulation results of thirteen SCMs from the baseline experiment. In this section we will investigate how sensitivity of these results to different forcing approaches. We will also show results from CCM3 with some improvements

on its deep convection scheme (ZM). It should be noted that not all SCMs participated in the sensitivity tests.

#### **a. Nudging approach**

Nudging approach specifies the total advective tendency from the observations and the simulated temperature and moisture are nudged toward the observations at each time step. Using the nudging approach can help reduce impacts of errors produced in the previous time step on simulations in the next time step.

With the nudging approach, the models that have problem in their triggering mechanisms, such as CCM3, Scripps and CSU, still significantly overestimate the observed precipitation rates during non-precipitation periods. The surface rainfall produced from these models show clear diurnal variations. This problem is not that significant in the rest of the models. This is consistent with the results seen in the baseline experiment.

#### **b. Forcing with a specified radiative heating rate**

This forcing method is identical to the baseline experiment except the radiative heating rate ( $Q_r$ ) is specified from the ECMWF analysis. This removes one more uncertainty in the multi-model intercomparison study. Here we only show the simulation results of precipitation, temperature and moisture for Subcase B (Figs. 18a-c). Temperature and moisture errors (Figs. 18b-c) are averaged between 115 – 915 hPa. It is seen that most of the models show very similar results as seen in the baseline experiment. It is noted that CCCma and PNNL/CCM2 show some sensitivity to this forcing approach. CCCma misses the first two precipitation events while it overestimates the last convective event when  $Q_r$  is fixed. In the baseline experiment, this model largely underestimates almost all the precipitation events. It also shows cold/moist biases in Figs. 18b-c in contrast to the warm/dry biases in Figs. 9b and 9e. By using the specified radiative heating rate, PNNL/CCM2 generally shows a warm bias instead of the cold bias in the baseline experiment. This may be because  $Q_r$  from the ECMWF analysis provides less radiative cooling than that from PNNL/CCM2 itself. In comparison with the specified  $Q_r$ , the radiative heating rates simulated from most models in the baseline experiment show more cooling in the upper and lower troposphere (not shown).

### c. Sensitivity tests to triggering mechanism and closure assumption

As mentioned before, among the models participating in the intercomparison, CCM3, CCM3/SIO and CCM3/SUNY are identical in every aspect except the parameterization of convection. Thus the differences among these models would give a measure of sensitivity of the simulations to convective parameterization. CCM3/SUNY uses a triggering mechanism proposed by Xie and M. Zhang (2000). They link the convective triggering mechanism with the large-scale dynamic forcing based on the observations to overcome the problem of using the CAPE only trigger in ZM. They assume that model convection occurs only when the large-scale dynamic forcing makes positive contributions to the existing positive CAPE. G. Zhang (2002) takes a further step to use the large-scale troposphere forcing (without the boundary layer forcing) to determine not only the timing, but also the amount of convection. He modified the CAPE closure for ZM so that the cloud base mass flux is determined by the CAPE generation rate due to the troposphere forcing. The new closure assumption is used in CCM3/SIO.

Figures 19a-c respectively show time series of the simulated precipitation rates and the averaged temperature and moisture biases between 115 – 915 hPa from CCM3, CCM3/SUNY and CCM3/SIO for Subcase B using the forcing approach in the baseline simulation. It is seen that both the revised ZM schemes generally well reproduce the observed precipitation rates, even for the first convective event of Subcase B, which almost all SCMs and CRMs fail to capture. This is because the overall large-scale forcing is strong enough to destabilize the atmosphere and make convection happen even though there is large-scale moisture divergence in the lower troposphere observed in this event (Fig. 3e). As a result, the large warm/dry biases shown in CCM3 are greatly reduced in CCM3/SUNY and CCM3/SIO. Compared to CCM3/SIO, CCM3/SUNY captures well the timing but overestimates the magnitude of convection. Therefore it produces larger errors in the temperature and moisture fields. This is because CCM3/SUNY still uses the same closure assumption as ZM uses to determine the cloud base mass flux, which is significantly reduced in the revised closure assumption in CCM3/SIO.

Cloud and radiation fields are also improved with the improved triggering mechanism and closure assumption. Figures 20a-b show the Taylor diagram for column cloud fraction and OLR at the TOA for Subcase B, respectively. In the figure, A1 denotes CCM3, M1 CCM3/SUNY and M2 CCM3/SIO. It is seen that both the revised ZM schemes improve the simulation of clouds

and OLR in terms of the correlation coefficient, the temporal variability and the RMS errors in comparison with the original ZM scheme.

## 6. Conclusion

Using the observations from the Summer ARM 1997 IOP and the data diagnosed from ten CRMs, fifteen SCMs have been evaluated under summertime midlatitude continental conditions with a focus on their cumulus parameterizations. The baseline experiment shows that most SCMs can reasonably reproduce the observed precipitation events that were well developed within the SCM domain. For those convective systems that occurred only within a small part of the domain, it has been shown that the domain averaged large-scale forcing may not properly capture the characteristics of these systems. This partly results in the failure of most models to simulate these precipitation events. It should be noted that one could not expect SCMs to be able to predict every single convective event right, given the current SCM domain size of  $370 \times 300 \text{ km}^2$ . However, the information of pdfs of thermodynamic properties in the boundary layer may help SCMs to get the statistics of convection occurrence right.

It has been shown that deficiencies associated with convective triggering mechanisms are mainly responsible for the large errors produced in some models. Convection in those models that use triggers based on the vertical integral of parcel buoyant energy is more active than those that use local parcel buoyancy if no additional constraints are used to control the onset of continental convection. The overactive convection leads to large systematic warm/dry biases in the troposphere in these models. It is also shown that a model with a non-penetrative convection scheme produces large systematic cold/moist biases in the troposphere due to underestimation of the depth of instability.

The radar estimated surface precipitation data show that stratiform precipitation has a considerable contribution (28%) to the total precipitation during the three precipitation periods chosen for this study. This is, however, not reproduced by SCMs; that is, the simulated surface precipitation is mainly from cumulus clouds for virtually all SCMs.

The updraft mass fluxes in the SCMs agree well with those in the CRMs. In contrast, most SCMs produce very weak downdraft mass fluxes compared with those diagnosed from the CRMs. This leads to larger net mass fluxes in the SCMs, especially in the lower and middle troposphere. As a result, most SCMs fail to produce the negative net mass fluxes in the lower

troposphere shown in the CRMs. Neglect of mesoscale circulations in the SCMs, uncertainties in the diagnosed mass fluxes from the CRMs and problems with cumulus parameterizations, such as the cloud base height and assumptions for downdrafts, are all responsible for the discrepancies. In the future, parameterization of mesoscale circulations needs to be incorporated into the models, in addition to further improvement of convective downdraft formulations (e.g. Donner 1993; Xu 1995). Reducing the uncertainties in the CRM mass fluxes, such as eliminating gravity wave contributions to the updrafts and downdrafts, is also needed so that CRMs can provide more reliable information for SCMs to improve their physical parameterizations.

It should be noted that deficiencies from other model parameterizations such as radiation, cloud and turbulent schemes could also affect the SCM simulations. From the heat and moisture budget analysis, however, this study has shown that convective heating and drying contribute mostly to the large intermodel differences in the troposphere. The intermodel differences from other physical processes are relatively small, compared to those from cumulus convection, except for the boundary layer. This may be related to the cases selected in this study that are mainly dominated by strong cumulus convection. It may also suggest that the interaction between convective and stratiform precipitation processes is not properly formulated in SCMs. Nevertheless, a detailed analysis of the interactions between cumulus convection and other physical processes should provide further guidance in the interpretation of model results.

Sensitivity tests have shown that similar results to the baseline experiment are obtained when different large-scale forcing approaches are used to drive the SCMs. It has also been shown that simulated results can be improved when deficiencies in some aspects of a cumulus parameterization are reduced.

## **Acknowledgments**

The authors would like to thank one of the anonymous referees for several valuable comments that helped to considerably improve the presentation of the paper. This research was supported primarily under the U. S. Department of Energy Atmospheric Radiation Measurement (ARM) Program. Work at LLNL was performed under the auspices of the U. S. Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48. Work at NASA Langley Research Center was partially supported by the NASA Earth Observation System (EOS) / Interdisciplinary Science Program under grant 291-01-91-10. Work at CSU was performed under ARM grant DE-FG03-95ER61968. Work for CCM3/SIO was supported by ARM grant DE-FG03-91ER61198. Work at PNNL was supported by the ARM Program; PNNL is operated for the DOE by Battelle Memorial Institute under contract DE-AC06-76RL01830. Work at SUNY Stony Brook was supported by ARM grant DE-FG02-98ER62570 and was also supported by NSF under grant ATM9701950. Work at the UK Met Office was supported by EU contract EVK2 CT199900051 (EUROCS). Work at the University of Utah was supported under ARM grant DE-FG03-94ER6179. Work at Scripps was supported in part by ARM grant DE-FG03-97-ER62338, by NOAA under grant NA77RJO453 and by NSF under grant ATM-9613764. Work at Dalhousie University was funded partly by NSERC. Work at CSIRO was funded partly through Australia's National Greenhouse Research Program.

## References

- Arakawa, A., and W. H. Schubert, 1974: Interaction of a cumulus cloud ensemble with the large-scale environment. Part I. *J. Atmos. Sci.*, **31**, 674-701
- Bechtold, P., et al. 2000: A GCSS model intercomparison for a tropical squall line observed during TOGA-COARE, II: Intercomparison of single-column models and a cloud-resolving model. *Q. J. R. Meteorol. Soc.*, **126**, 8665-888
- Bechtold, P., E. Bazile, F. Guichard, P. Mascart, and E. Richard, 2001: A mass flux convection scheme for region and global models. *Q. J. R. Meteorol. Soc.*, **127**, 869-886
- Browning, K. A., 1994: GEWEX cloud system study (GCSS) science plan. *IGPO Publication Series*, **No. 11**, International GEWEX Project Office, 84pp
- Cederwall, R. T., S. K. Krueger, S. Xie, and J. Yio, 2000: ARM/GCSS Single Column Model (SCM) Intercomparison, Procedures for Case 3: Summer 1997 SCM IOP. *LLNL Report*, UCRL-ID-141823, 20pp
- Cheng, M.-D., and M. Yanai, 1989: Effects of downdrafts and mesoscale convective organization on the heat and moisture budgets of tropical cloud clusters. Part III: Effects of mesoscale convective organization. *J. Atmos. Sci.*, **46**, 1566-1588
- Del Genio, A. D., and M.-S. Yao, 1993: Efficient cumulus parameterization for long-term climate studies: The GISS scheme in *The Representation of Cumulus Convection in Numerical Models*, *Meteorol. Monogr.*, **46**, edited by K. Emanuel and D. Raymond, 246pp., Am. Meteorol. Soc., Boston, Mass.
- Del Genio, A. D., et al., 1996: A prognostic cloud water parameterization for global climate models. *J. Climate*, **9**, 270-304
- Ding, P., and D. A. Randall, 1998: A cumulus parameterization with multiple cloud base levels. *J. Geophys. Res.*, **103**, 11341-11354
- Donner, L. J., 1993: A cumulus parameterization including mass fluxes, vertical momentum dynamics, and mesoscale effects. *J. Atmos. Sci.*, **50**, 889-906
- Ghan, S. J., L. R. Leung, and Q. Hu, 1997: Application of cloud microphysics to NCAR CCM2. *J. Geophys. Res.*, **102**, 16,507-16,527.
- Ghan, S. J., et al., 2000: An intercomparison of single column model simulations of summertime midlatitude continental convection. *J. Geophys. Res.*, **105**, 2091-2124.

- Gregory, D., 2001: Estimation of entrainment rate in simple models of convective clouds. *Q. J. R. Meteorol. Soc.*, **127**, 53-72
- Gregory, D., and F. Guichard, 2001: Aspects of the parameterization of organized convection: contrasting cloud resolving model and single column realizations, submitted to *Q. J. R. Meteorol. Soc.*
- Gregory, D., J.-J. Morcrette, C. Jakob, A. M. Beljaars, and T. Stockdale, 2000: Revision of convection, radiation and cloud schemes in the ECMWF Integrated Forecasting System. *Q. J. R. Meteorol. Soc.*, **126**, 1686-1710
- Gregory, D., 1995: Convection scheme, part 1 of UM Documentation Paper 27, NWP-Div., Meteorological Office, London Road, Bracknell, Berks. RG12 2SZ
- Gregory, D., and P. R. Rowntree, 1990: A mass flux convection scheme with representation of cloud ensemble characteristics and stability-dependent closure. *Mon. Weather Rev.*, **118**, 1483-1506
- Grell, G. A., Y. -H. Kuo, and R. Pasch, 1991: Semiprognostic tests of cumulus parameterization schemes in the middle latitudes. *Mon. Weather Rev.*, **119**, 5-31
- Hack, J. J., B. A. Boville, B. P. Briegleb, J. T. Kiehl, P. J. Rasch, and D. L. Williamson, 1993: Description of the NCAR community climate model (CCM2), *NCAR Tech. Note, NCAR TN-336-STR*, National Center for Atmospheric Research, Boulder, CO, 108pp
- Hack, J. J., 1994: Parameterization of moist convection in the National Center for Atmospheric Research community climate model (CCM2). *J. Geophys. Res.*, **99**, 5551-5568
- Hack, J. J., J. A. Pedretti, and J. C. Petch, 1998: SCCM User's Guide (<http://www.cgd.ucar.edu/cms/sccm/sccm.html>).
- Houze, R. A., and C.-P. Cheng, 1981: Inclusion of mesoscale updrafts and downdrafts in computations of vertical fluxes by ensembles of tropical clouds. *J. Atmos. Sci.*, **38**, 1751-1770
- Iacobellis S. F., and R. J. C. Somerville, 2000: Implication of microphysics for cloud-radiation parameterizations: Lessons from TOGA COARE. *J. Atmos. Sci.*, **57**, 161-183
- Johnson, R. H., 1980: Diagnosis of convective and mesoscale motions during Phase III of GATE. *J. Atmos. Sci.*, **37**, 733-753



- Johnson, R. H., and P. J. Hamilton 1980: The relationship of surface pressure features to the precipitation and airflow structure of an intense midlatitude squall line. *Mon. Wea. Rev.*, **116**, 1444-1472
- McFarlane N. A., G. J. Boer, J-P. Blanchet, and M. Lazare, 1992: The Canadian Climate Center second-generation general circulation model and its equilibrium climate. *J. Climate*, **5**, 1013-1044
- Kiehl, J. T., B. Boville, B. Briegleb, J. Hack, P. Rasch, and D. Williamson, 1996: Description of the NCAR Community Climate Model (CCM3). *NCAR Technical Note, NCAR/TN-420+STR*, 151pp
- Krueger, S. K., et al., 2002: Intercmparison of multi-day simulations of convection during TOGA COARE with several cloud-resolving and single-column models. (to be submitted)
- Lafore, J. P. et al., 1998: The Meso-NH Atmospheric Simulation System. Part I: Adiabatic formulation and control simulations. *Annales Geophysicae*, **16**, 90-109
- Lohmann, U., N. McFarlane, L. Levkov, K. Abdella, and F. Albers, 1999: Comparing different cloud schemes of a single column model by using mesoscale forcing and nudging technique. *J. Climate*, **12**, 438-461
- Lord, S. J., and A. Arakawa, 1982: Interaction of a cumulus cloud ensemble with the large-scale environment. Part III: Semiprognostic test of Arakawa-Schubert cumulus parameterization. *J. Atmos. Sci.*, **39**, 88-103
- Moorthi, S., and M. J. Suarez, 1992: Relaxed Arakawa-Schubert, A parameterization of moist convection for general circulation models. *Mon. Weather Rev.*, **120**, 978-1002
- Moran, K. P., B. E. Martner, M. J. Post, R. A. Kropfli, D. C. Welsh, and K. B. Widerner, 1998: An attended cloud-profiling radar for use in climate research. *Bull. Amer. Meteor. Soc.*, **79**, 443-455
- Ogura, Y., and H. –Y. Jiang, 1985: A modeling study of heating and drying effects of convective clouds in an extratropical mesoscale convective system. *J. Atmos. Sci.*, **42**, 2478-2492
- Pan, D. –M., and D. A. Randall, 1998: A cumulus parameterization with a prognostic closure. *Q. J. R. Meteorol. Soc.* **125**, 949-982
- Pope, V. D., M. Gallani, P. R. Rowntree, and R. A. Stratton, 2000: The impact of new physical parametrizations in the Hadley Centre climate model – HadAM3. *Climate Dyn.*, **16**, 123-146

- Randall, D. A., and D. G. Cripe, 1999: Alternative methods for specification of observed forcing in single-column models and cloud system models. *J. Geophys. Res.*, **104**, 24527-24545
- Randall, D. A., K-M. Xu, R. J. C. Somerville and S. Iacobellis, 1996: Single-column models and cloud ensemble models as links between observations and climate models. *J. Climate*, **9**, 1683-1697
- Raymond D. J., 1993: Observational constraints on cumulus parameterization in *The Representation of Cumulus Convection in Numerical Models*, Meteorol. Monogr., **46**, edited by K. Emanuel and D. Raymond, 246pp., Am. Meteorol. Soc., Boston, Mass.
- Redelsperger, J.-L., et al., 2000: A GCSS model intercomparison for a tropical squall line observed during TOGA-COARE. Part I: CRM results. *Q. J. R. Meteorol. Soc.* **126**, 823-863
- Rotstayn, L. D., 1997: A physical based scheme for the treatment of stratiform clouds and precipitation in large-scale models. I: Description and evaluation of the microphysical processes. *Q. J. R. Meteorol. Soc.* **123**, 1227-1282
- Rotstayn, L. D., B. F. Ryan, and J. J. Katzfey, 2000: A scheme for calculation of the liquid fraction in mixed-phase stratiform clouds in large-scale models. *Mon. Weather Rev.*, **128**, 1070-1088
- Stokes, G. M., and S. E. Schwartz, 1994: The Atmospheric Radiation Measurement (ARM) program: Programmatic background and design of the cloud and radiation test bed. *Bull. Amer. Meteor. Soc.*, **75**, 1202-1221
- Sud, Y. C., and G. K. Walker, 1993: A rain-evaporation and downdraft parameterization to complement a cumulus updraft scheme and its evaluation using GATE data. *Mon. Weather Rev.*, **121**, 3019-3039
- Sud, Y. C., and G. K. Walker, 1999a: Microphysics of clouds with the relaxed Arakawa-Schubert scheme (McRAS). Part I, Design and evaluation with GATE phase III data. *J. Atmos. Sci.*, **56**, 3196-3220
- Sud, Y. C., and G. K. Walker, 1999b: Microphysics of clouds with the relaxed Arakawa-Schubert scheme (McRAS). Part II, Implementation and Performance in GEOS II GCM. *J. Atmos. Sci.*, **56**, 3221-3240
- Takacs, L. L., A. Molod, and T. Weng, 1994: Goddard Earth Observing System (GEOS) general circulation model (GCM) version 1. *NASA Tech. Memo. 104606*, Vol. 1, 97 pp.

- Taylor, K. E., 2000: Summarizing multiple aspects of model performance in a single diagram. *J. Geophys. Res.*, **106**, 7183-7192
- Tiedtke, M., 1989: A comprehensive mass flux scheme for cumulus parameterization in large-scale models. *Mon. Weather Rev.*, **117**, 1779-1800
- Thompson, R. M., Jr., S. W., Payne, E. E. Recker, and R. J. Reed, 1979: Structure and properties of synoptic-scale wave disturbances in the introtropical convergence zone of the eastern Atlantic. *J. Atmos. Sci.*, **36**, 53-72
- von Salzen, K., and N. A. McFarlane, 2002: Parameterization of the effects of lateral and cloud-top entrainment in transient shallow cumulus clouds. *J. Atmos. Sci.*, in press
- Wu, X., 1993: Effects of cumulus ensembles and mesoscale stratiform clouds in midlatitude convective systems. *J. Atmos. Sci.*, **50**, 2496-2518
- Webster, P. J., and R. Lukas, 1992: TOGA COARE: The coupled ocean-atmosphere response experiment. *Bull. Amer. Meteor. Soc.*, **73**, 1377-1417
- Xie, S. C., and M. H. Zhang, 2000: Impact of the convective triggering function on single-column model simulations. *J. Geophys. Res.*, **105**, 14983-14996
- Xu, K.-M., and A. Arakawa, 1992: Semi-prognostic tests of the Arakawa-Schubert cumulus parameterization using simulated data. *J. Atmos. Sci.*, **49**, 2421-2436
- Xu, K.-M., 1995: Partitioning mass, heat and moisture budgets of explicitly simulated cumulus ensembles into convective and stratiform components. *J. Atmos. Sci.*, **52**, 551-573
- Xu, K. -M., and D. A. Randall, 2000: Explicit simulation of midlatitude cumulus ensembles: comparison with ARM data. *J. Atmos. Sci.*, **57**, 2839-2858
- Xu, K. -M., et al., 2002: An intercomparison of cloud-resolving models with the ARM summer 1997 IOP data. *Q. J. R. Meteorol. Soc.* (to appear).
- Yanai, M., S. Esbensen, and J. Chu, 1973: Determination of bulk properties of tropical cloud clusters from large-scale heat and moisture budgets. *J. Atmos. Sci.*, **30**, 611-627
- Yanai, M., J. Chu, T. E. Stark, and T. Nitta, 1976: Response of deep and shallow tropical maritime cumuli to large-scale processes. *J. Atmos. Sci.*, **33**, 976-991
- Zhang, G. J., and N. A. McFarlane, 1995: Sensitivity of climate simulations to the parameterization of cumulus convection in the Canadian Climate Center general circulation model. *Atmosphere-Ocean*, **33**, 407-446

- Zhang, G. J., 2002: Convective quasi-equilibrium in midlatitude continental environment and its effect on convection parameterization. (Accepted by *J. Geophys. Res.*).
- Zhang, M. H., and J. L. Lin, 1997: Constrained variational analysis of sounding data bases on column-integrated budgets of mass, heat, moisture, and momentum: Approach and application to ARM measurements. *J. Atmos. Sci.*, **54**, 1503-1524
- Zhang, M. H., J. L. Lin, R. T. Cederwall, J. J. Yio, and S. C. Xie, 2001: Objective analysis of ARM IOP Data: Method and sensitivity. *Mon. Weather Rev.*, **129**, 295-311

### Figure caption

- Figure 1. Time series of the observed surface precipitation rates during Summer 1997 IOP. Labels A, B and C in the plot are the three subcases selected in the study.
- Figure 2. Satellite infrared cloud images corresponding to the peaks of the eight strong precipitation events that occurred on Days 179, 180-181, 190, 191, 192, 195, 196 and 197-198 during Summer 1997 IOP.
- Figure 3. Time-height cross section of observed large-scale advective tendencies of potential temperature (a – c;  $\text{K day}^{-1}$ ) and moisture (d – e;  $\text{g kg}^{-1} \text{ day}^{-1}$ ) and surface precipitation rates (f – h;  $\text{mm day}^{-1}$ ) for Subcases A, B and C.
- Figure 4. Time series of the observed (black line) and simulated total ((a) – (c)) and stratiform ((d) – (f)) surface precipitation rates for Subcases A, B and C.
- Figure 5. A Taylor diagram showing how closely the simulated surface precipitation rates match with the observation for total precipitation rate (a) and convective precipitation rate (b) for Subcase B. This polar style graph is designed to show the correlation coefficient for a field between observations and simulations (azimuthal position), the standard deviation of observed and simulated fields (radial distances), and the root-mean square (RMS) errors once any overall bias has been removed (the distance connecting the observation and simulation points). Note that the RMS error and the standard deviation are normalized by the standard deviation of the corresponding observed field. The standard deviation of the observed field is normalized by itself and it therefore always be plotted at unit distance from the origin along the abscissa. The distance connecting the observation and simulation points represents the root-mean square (RMS) errors. A1 denotes CCM3, A2 Scripps, A3 CCCma1\_SP, B1 CCCma, B2 CSU, B3 McRAS, B4 GFDL, C1 GISS, C2 CSIRO, C3 MOUM, C4 PNNL/CCM2, C5 MesoNH, X ten CRMs consensus and Obs observations.
- Figure 6. Vertical profiles of moist static energy (MSE, solid) and saturated MSE (dashed) corresponding to the beginning of four selected precipitation events that occurred on Days 180-181, 190, 191 and 195. (a) – (d) MSE is calculated from the sounding profiles

at the station where convection is occurring. (e) – (h) MSE is calculated from the ARM SCM domain mean soundings.

Figure 7. Time series of (a) CAPE and precipitation and (b) CIN and precipitation for the last ten days of Summer 1997 IOP, which covers the period of Subcases B and C.

Figure 8. Same as Fig. 5 except for (a) column cloud fractions and (b) TOA outgoing longwave radiation fluxes and that the standard deviations are not normalized.

Figure 9. Time evolution of the column mean (between pressure levels 115 – 915 hPa) temperature ((a) – (c)) and moisture ((d) – (f)) errors produced by the SCMs for Subcase A, B and C.

Figure 10. Vertical profiles of time average temperature ((a) – (c)) and moisture ((d) – (f)) errors produced by the SCMs over Subcases A, B and C.

Figure 11. Vertical profiles of updraft, downdraft and net cloud mass fluxes averaged over precipitation periods (observed precipitation rates  $\geq 0.36 \text{ mm day}^{-1}$ ) and non-precipitation periods (observed precipitation rates  $< 0.36 \text{ mm day}^{-1}$ ) of Subcases A, B and C. Black solid line is the observed large-scale mean mass flux ( $-\bar{w}$ ). Black dashed line and small bar represent the mean of the mass fluxes across the ten CRMs and the standard deviations from the mean, respectively. (a) – (c) respectively represent the updrafts, downdrafts and net cloud mass fluxes during precipitation periods. (d) – (f) are the same as (a) – (c) except for non-precipitation periods.

Figure 12. Vertical profiles of (a) updraft, (b) downdraft and (c) net cumulus mass fluxes estimated from UCLA/CSU CRM averaged over precipitation periods (observed precipitation rates  $\geq 0.36 \text{ mm day}^{-1}$ ) of Subcases A, B and C. Solid lines represent the total mass fluxes, dotted lines convective-scale contributions and dashed lines mesoscale contributions.

Figure 13. Apparent heat sources ( $Q_1$ ) and contributions from cumulus convection ( $Q_{1c}$ ), radiation ( $Q_r$ ) and turbulent processes ( $Q_{1v}$ ), averaged over precipitation periods of Subcases A, B and C ( $\text{K day}^{-1}$ ).

Figure 14. Same as Fig. 13 except for non-precipitation periods of Subcases A, B and C.

Figure 15. Apparent moisture sinks ( $Q_2$ ) and contributions from cumulus convection ( $Q_{2c}$ ) and turbulent processes ( $Q_{2v}$ ), averaged over precipitation periods of Subcases A, B and C ( $\text{k day}^{-1}$ ).


Figure 16. Same as Fig. 15 except for the non-precipitation periods of Subcases A, B and C.

Figure 17. Time series of the observed and simulated surface total precipitation rates ( $\text{mm day}^{-1}$ ) using the nudging approach for Subcases A, B and C.

Figure 18. Time series of the observed and simulated surface total precipitation rates ( $\text{mm day}^{-1}$ ) and the column mean (between pressure levels 115 – 915 hPa) temperature and moisture errors produced by the SCMs for Subcase B, using the forcing with a specified radiative heating rate. (a) Surface precipitation rates. (b) Temperature. (c) Moisture.

Figure 19. Time series of the observed and simulated total precipitation rates ( $\text{mm day}^{-1}$ ) and the column mean (between pressure levels 115 – 915 hPa) temperature and moisture errors produced by the SCMs for Subcase B, using the same forcing approach as in the baseline experiment. Thick solid line is for the observations, thin solid lines are for CCM3, dotted lines are for CCM3/SUNY and dashed lines are for CCM3/SIO. (a) Surface precipitation rates. (b) Temperature. (c) Moisture.

Figure 20. A Taylor diagram to compare (a) column cloud fraction and (b) OLR at the TOA produced from CCM3, CCM3/SUNY and CCM3/SIO. A1 denotes CCM3, M1 CCM3/SUNY and M2 CCM3/SIO. See Fig. 5 for a detailed explanation.





## GOES Cloud Images

